

ON HÖLDER CONTINUOUS RIEMANNIAN AND FINSLER METRICS

ALEXANDER LYTCHAK AND ASLI YAMAN

ABSTRACT. We discuss smoothness of geodesics in Riemannian and Finsler metrics.

1. INTRODUCTION

In this paper we improve and generalize a result of Calabi and Hartmann in [CH70]. They study smoothness of isometries between equal-dimensional manifolds with α -Hölder continuous Riemannian metrics and prove that each isometry is of class $\mathcal{C}^{1,\alpha}$. The main tool in the proof is Theorem 3.1 of [CH70], stating that geodesics in α -Hölder Riemannian metrics are uniformly $\mathcal{C}^{1,\alpha}$, but the proof of this theorem is not correct. Although the result on the smoothness of isometries is true, as Taylor has shown by a different method in [Tay], Theorem 3.1 is wrong for $\alpha < 1$.

Indeed one of our results is the following:

Theorem 1.1. *For each $0 < \alpha \leq 1$ set $\beta = \frac{\alpha}{2-\alpha}$. There is an α -Hölder Riemannian metric on \mathbb{R}^2 , such that geodesics near the origin are not uniformly $\mathcal{C}^{1,l}$ for any $l > \beta$.*

Actually it seems to be possible but technically not trivial to construct an α -Hölder continuous Riemannian metric for which some geodesic γ is not $\mathcal{C}^{1,l}$ for all $l > \beta$. This would show that there are manifolds (of different dimensions) with α -Hölder Riemannian metrics and an distance-preserving embedding of one of them into another that is at best $\mathcal{C}^{1,\beta}$, in contrast to the $\mathcal{C}^{1,\alpha}$ smoothness of isometries shown by Taylor.

Remark 1.1. If α goes to 1 then the regularity of geodesics is almost $\mathcal{C}^{1,\alpha}$, whereas if α goes 0 one only gets a bit more than half of the expected regularity!

In contrast to this result the first author proved in [Lyt] that geodesics in $\mathcal{C}^{1,\alpha}$ submanifolds of smooth Riemannian manifolds are uniformly $\mathcal{C}^{1,\alpha}$. This gives us the following non-embeddability result:

1991 *Mathematics Subject Classification.* 53B40, 53B20.

Key words and phrases. Geodesics, isometric embeddings.

Corollary 1.2. *For $0 < \alpha < 1$ and $\beta = \frac{\alpha}{2-\alpha}$ there are α -Hölder continuous Riemannian metrics that (even locally) do not admit $\mathcal{C}^{1,l}$ arcwise isometric embeddings into any smooth Riemannian manifold, for any $l > \beta$.*

This result should be compared to the theorems of Nash, stating that each manifold with a continuous Riemannian metric has a \mathcal{C}^1 arcwise isometric embedding into some \mathbb{R}^n and that for $k \geq 3$ each Riemannian manifold with a \mathcal{C}^k -Riemannian metric has a \mathcal{C}^k arcwise isometric embedding into some Euclidean space.

Remark 1.2. For $\alpha = 1$ a statement similar to Corollary 1.2 still holds, but for another reason. Namely it is shown in [Lyt], that each $\mathcal{C}^{1,1}$ submanifold of a smooth Riemannian manifold has curvature bounded from both sides with respect to the inner metric, whereas it is certainly not true even for a general Riemannian metric of class $\mathcal{C}^{1,l}$, for any fixed $l < 1$.

The proof of Theorem 3.1 in [CH70] only shows that geodesics in α -Hölder Riemannian metrics are uniformly $\mathcal{C}^{1,\frac{\alpha}{2}}$. We repeat their argument and generalize this result to sufficiently convex Finsler metrics (see Definition 2.3 and Definition 3.1 for precise definitions) :

Theorem 1.3. *Let $(M, |\cdot|)$ be a manifold with an α -Hölder Finsler structure $|\cdot|$, such that the norms $|\cdot|_x$ are locally uniformly of convexity type p . Then geodesics with respect to the Finsler metric in M are locally uniformly $\mathcal{C}^{1,\frac{\alpha}{p}}$.*

Exactly as in the last part of [CH70], Theorem 1.3 implies that every distance-preserving embedding between two manifolds as in Theorem 1.3 is $\mathcal{C}^{1,\frac{\alpha}{p}}$.

Finally we prove the main result of this paper, showing that Theorem 1.1 is optimal:

Theorem 1.4. *Let $(M, \langle \cdot, \cdot \rangle_x)$ be a manifold with an α -Hölder Riemannian metric. Then geodesics in M are locally uniformly $\mathcal{C}^{1,\beta}$ for $\beta = \frac{\alpha}{2-\alpha}$.*

After recalling some basic facts about $\mathcal{C}^{1,\alpha}$ maps we study curves in a manifold M with a Finsler structure and compare the lengths of curves to their lengths measured in some tangent space. Using these estimations we prove Theorem 1.3 and Theorem 1.4. The main idea is that a curve $\gamma : [a, b] \rightarrow M$ that is not $\mathcal{C}^{1,l}$ admits for arbitrary $A > 0$ arbitrary closed points $t, s \in [a, b]$ such that $\gamma(\frac{t+s}{2})$ is at least $A||t - s||^l$ far away from the straight line η connecting $\gamma(t)$ and $\gamma(s)$. Considering on M a Banach space metric induced by a norm $|\cdot|_x$ for each x , we see from the fact that the norms are sufficiently convex, that the length of γ is much bigger than the length of η with respect to this norm. Now using the fact that the lengths of η and γ with respect to this norm do not differ to much from their actual lengths

in M one get a contradiction if γ is a geodesic. The idea of Hartman and Calabi in order to prove Theorem 1.3 was to compute the lengths of η and γ relative to a tangent space at the fixed point $\gamma(t_0)$. A better estimation providing Theorem 1.4 is achieved by comparing at each t the quantities $|\gamma'(t)|_{\gamma(t)}$ and $|\eta'(t)|_{\eta(t)}$.

After preliminaries in Section 2, we study in Section 3 Hölder continuous Finsler structures, discuss examples and basics of the structure and recall the comparisons of Calabi and Hartmann of lengths of curves measured in M with their lengths with respect to the constant Finsler structure $|\cdot|_x = |\cdot|_{x_0}$. In Section 4 we prove Theorem 1.3 and Theorem 1.4. Finally in Section 5 we prove Theorem 1.1 by giving a counterexample to the result of Calabi and Hartman.

We are grateful to Werner Ballmann and Juan Souto for helpful suggestions. We are indebted to Reiner Schätzle for indicating a mistake in the previous version of our counterexample. We are also thankful to Jonathan Wahl for bringing the work of Michael Taylor to our attention.

2. PRELIMINARIES

Let us start with some basic conventions that shall be followed throughout this work. A geodesic in a metric space X is an arclength parametrised curve $\gamma : [a, b] \rightarrow X$ satisfying $d(\gamma(a), \gamma(b)) = b - a$. By $\|\cdot\|$ we will denote the Euclidean norm in Euclidean spaces. Given a \mathcal{C}^t manifold M a chart is a \mathcal{C}^t diffeomorphism which sends an open subset of \mathbb{R}^n onto an open subset of M . We shall always omit the neighbourhood in M and identify a chart with its image in \mathbb{R}^n .

2.1 Hölder maps

A map $f : (M, d_1) \rightarrow (N, d_2)$ between metric spaces M and N is (C, α) -Hölder, for $C > 0, 1 \geq \alpha > 0$ if for all $x, z \in X$ one has $d_2(f(x), f(z)) \leq C d_1(x - z)^\alpha$. The map is called locally α -Hölder if M is covered by open subsets U_j , such that each restriction $f : U_j \rightarrow N$ is (C_j, α) -Hölder for some $C_j = C(U_j)$.

Let M and N be locally compact spaces. Let \mathcal{F} be a family of maps $f_a : D_a \rightarrow N$ defined on subsets D_a of M . For subsets $X \subset M$ and $Y \subset N$ denote by $\mathcal{F}_{X,Y}$ the subfamily of all maps $f \in \mathcal{F}$ satisfying $f_a(X \cap D_a) \subset Y$.

Definition 2.1. *We will say that the family \mathcal{F} is locally uniformly α -Hölder, if for all compact subsets $X \subset M$ and $Y \subset N$ there is a positive $C = C(X, Y) > 0$, such that the restriction $f_a : (D_a \cap X) \rightarrow Y$ is (C, α) -Hölder for all $f_a \in \mathcal{F}_{X,Y}$.*

Since M and N are by definition locally compact, the family \mathcal{F} is locally uniformly α -Hölder if and only if the condition of Definition 2.1 is satisfied

for all open relatively compact subsets $X = U_j$ and $Y = V_i$ of a covering U_j of M resp. V_i of N .

2.2 $\mathcal{C}^{1,\alpha}$ -maps

A map $f : U \rightarrow \mathbb{R}^m$ defined on an open subset U of \mathbb{R}^n is locally $\mathcal{C}^{1,\alpha}$ if it is \mathcal{C}^1 and its differential $Df : U \rightarrow \mathbb{L}(\mathbb{R}^n, \mathbb{R}^m)$ is locally α -Hölder where $1 \geq \alpha > 0$.

The following lemma is in fact a consequence of a more general Lemma 2.2. However we state the lemma and give a proof for the later reference in the case of curves.

Lemma 2.1. *Let $\gamma : [a, b] \rightarrow \mathbb{R}^n$ be a \mathcal{C}^1 curve with $\|\gamma'(t) - \gamma'(s)\| \leq C\|s - t\|^\alpha$ for all $s, t \in (a, b)$. Then for all r with $a < r < b$ we have $\|\gamma'(r)\|b - a\| - (\gamma(b) - \gamma(a))\| \leq C\|b - a\|^{1+\alpha}$.*

Proof. We have $\gamma(b) - \gamma(a) = \int_a^b \gamma'(x)dx = \int_a^b \gamma'(r)dx + \int_a^b (\gamma'(x) - \gamma'(r))dx$. Using $\|\gamma'(x) - \gamma'(r)\| \leq C\|b - a\|^\alpha$ we obtain the result. \square

In particular this shows that under the conditions of Lemma 2.1 the distance between $\gamma(r)$ and the affine line through $\gamma(a)$ and $\gamma(b)$ is at most $C\|b - a\|^{1+\alpha}$.

Definition 2.2. *A family \mathcal{F} of maps $f_a : U_a \rightarrow \mathbb{R}^m$ defined on open subsets U_a of \mathbb{R}^n is called locally uniformly $\mathcal{C}^{1,\alpha}$ if the family is locally uniformly Lipschitz and the family $\tilde{\mathcal{F}}$ of differentials $\tilde{f}_a = (f_a, Df_a) : U_a \rightarrow \mathbb{R}^m \times \mathbb{L}(\mathbb{R}^n, \mathbb{R}^m)$ is locally uniformly α -Hölder.*

We will need the following characterisation of locally uniformly $\mathcal{C}^{1,\alpha}$ families. That result is given as Lemma 2.1 in [CH70].

Lemma 2.2. *A family \mathcal{F} of locally uniformly Lipschitz maps $f_a : U_a \rightarrow \mathbb{R}^m$ defined on open subsets U_a of \mathbb{R}^n is locally uniformly $\mathcal{C}^{1,\alpha}$, if for each ball $B \subset \mathbb{R}^m$, one can cover U_a by open subsets O_j , such that for each all $f \in \mathcal{F}_{O_j, B}$ all $x \in O_j$ and all $h \in \mathbb{R}^n$ such that $x + h$ and $x - h$ are in O_j one has $\|f(x + h) + f(x - h) - 2f(x)\| \leq C\|h\|^{1+\alpha}$ for some constant $C = C(O_j)$.*

Since the composition of $\mathcal{C}^{1,\alpha}$ maps is again $\mathcal{C}^{1,\alpha}$ the notion of a $\mathcal{C}^{1,\alpha}$ manifold is well defined. Moreover one can speak about locally uniformly $\mathcal{C}^{1,\alpha}$ families of maps between two $\mathcal{C}^{1,\alpha}$ manifolds.

2.3 Norms of convexity type p

Let $(V, |\cdot|)$ be a (finite dimensional) normed vector space. Recall that the modulus of convexity $\delta = \delta_{|\cdot|}$ of the norm is defined by $\delta(\epsilon) = \inf\{1 - \frac{|v+w|}{2} \mid |v|, |w| \leq 1, |v - w| \geq \epsilon\}$ (See [LT79] p. 59 - 66 for further discussion on this).

Definition 2.3. Let $\kappa > 0, p \geq 2$ be given. We will say that the norm $|\cdot|$ is κ -convex of type p , if $\delta(\epsilon) \geq \kappa\epsilon^p$ for all $\epsilon < \kappa$.

Remark 2.1. Informally speaking, the above definition means that V is at least as convex as the Lebesgue space \mathcal{L}^p (see [LT79] p.63).

Example 2.2. If $(V, |\cdot|)$ is a κ -convex space of type p , then the same holds for each subspace of V . On the other hand $(V, |\cdot|)$ is κ -convex of type p if this is true for each two-dimensional subspace W of V .

Example 2.3. The Euclidean space is κ -convex of type 2 with $\kappa = \frac{1}{10}$. More generally the statement that the norm on V is κ -convex of type 2 is equivalent to the fact that the second fundamental form of the unit sphere S of V is bounded below by a constant $\epsilon = \epsilon(\kappa) > 0$.

We can now reformulate the definition in a way we will use it:

Lemma 2.3. Let the norm $|\cdot|$ on the vector space V be κ -convex of type p . Let $K, \alpha > 0$. Then there are positive $\epsilon = \epsilon(\kappa, K, \alpha)$ and $\lambda = \lambda(\kappa, K, \alpha)$ such that for all numbers h with $h \leq \epsilon$ and all $v, w \in V$ with $|v|, |w| \leq h + Kh^{1+\alpha}$ and $|v + w| \geq 2h - Kh^{1+\alpha}$ one has $|v - w| \leq \lambda h^{1+\frac{\alpha}{p}}$.

Proof. First we note that if $|\cdot|$ is κ -convex of type p then for all $H \leq \kappa^{p+1} = \epsilon'$ and for all $|v_0|, |w_0| \leq 1$ the inequality $2 - |v_0 + w_0| \leq H$ implies $|v_0 - w_0| \leq \lambda' H^{\frac{1}{p}}$ where $\lambda' = \kappa^{-\frac{1}{p}}$.

Now consider $v_0 = \frac{v}{h+Kh^{1+\alpha}}$ and $w_0 = \frac{w}{h+Kh^{1+\alpha}}$. Then $|v_0|, |w_0| \leq 1$ and $2 - |v_0 + w_0| \leq 3Kh^\alpha$. Thus for h with $3Kh^\alpha \leq \epsilon'$ the above remark implies that $|v_0 - w_0| \leq \lambda'(3Kh^\alpha)^{\frac{1}{p}} = (\frac{3K}{\kappa})^{\frac{1}{p}} h^{\frac{\alpha}{p}}$. Thus by setting $\epsilon = (\frac{\epsilon'}{3K})^{\frac{1}{\alpha}}$ and $\lambda = (\frac{3K}{\kappa})^{\frac{1}{p}}$, we have the result. \square

3. FINSLER STRUCTURES

3.1 Definition and examples

A Finsler structure on a \mathcal{C}^1 -manifold M is a continuous map $|\cdot| : TM \rightarrow [0, \infty)$ on the tangent bundle TM such that for all $x \in M$ the restriction $|\cdot|_x : T_x M \rightarrow [0, \infty)$ is a norm. Given a Finsler structure one can speak about lengths of locally Lipschitz curves $\gamma : [a, b] \rightarrow M$ given by $L(\gamma) = \int_a^b |\gamma'(t)|_{\gamma(t)} dt$. This defines as usual a length metric d on M which is called the Finsler metric. Remark that the Finsler metrics defined by two different Finsler structures are locally bi-Lipschitz equivalent. In particular the Finsler metric defines the usual locally compact topology on M . Thus by the theorem of Arzela-Ascoli M is covered by open sets U_j , such that each two points $x, z \in U_j$ are connected in M by a geodesic.

Definition 3.1. A Finsler structure $|\cdot|$ on a manifold M is locally uniformly of type p , if for each compact subset X of M there is a constant $\kappa = \kappa(X) > 0$ such that, for each $x \in X$ the norm $|\cdot|_x$ is κ -convex of type p .

For an immersion $\phi : N \rightarrow M$ the Finsler structure on M defines a Finsler structure on N through $|v|_x = |D_x\phi(v)|_{\phi(x)}$. In particular we get a Finsler structure on each \mathcal{C}^1 -submanifold of M and on each chart $U \subset \mathbb{R}^n$ of M .

On an open subset U of \mathbb{R}^n with a Finsler structure $|\cdot|$ we consider the function $\tilde{o} : U \times U \rightarrow \mathbb{R}^+$ given by $\tilde{o}(x, y) = \sup_{\|v\| \leq 1} ||v|_x - |v|_y||$. Recall that $\|\cdot\|$ denotes the Euclidean norm. The function \tilde{o} is continuous and vanishes on the diagonal. Hence on each relatively compact open subset V of U the function $o_V(r) = \sup\{\tilde{o}(x, y) \mid x, y \in V, \|x - y\| \leq r\}$ is finite, continuous, non-decreasing and satisfies $o_V(0) = 0$. Moreover since V is relatively compact we can find a constant $C > 0$ such that $\frac{1}{C}|v|_x \leq \|v\| \leq C|v|_x$ for all $v \in \mathbb{R}^n$ and all $x \in V$.

Let W be an open subset of \mathbb{R}^m and let $\phi : W \rightarrow V$ be an immersion, satisfying $\|\phi(x) - \phi(y)\| \leq L\|x - y\|$ for some constant $L \geq 0$. Consider the quantity $u(r) = \sup_{\|v\| \leq 1, \|x - y\| \leq r} \|D_x\phi(v) - D_y\phi(v)\|$. Then for the induced Finsler structure on W it is easily checked that $o_W(r) \leq Lo_V(Lr) + C \cdot u(r)$.

Hence if for all r we have $o_V(r) \leq Cr^\alpha$ for some $C \geq 0$ and $1 \geq \alpha > 0$, and if the map ϕ is $\mathcal{C}^{1,\alpha}$, we obtain $o_W(r) \leq C'r^\alpha$ for some $C' \geq 0$. Hence the following definition is meaningful:

Definition 3.2. A Finsler structure on a $\mathcal{C}^{1,\alpha}$ -manifold M is called a \mathcal{C}^α -Finsler structure if for each chart U and for each relatively compact open subset V of U one has $o_V(r) \leq Cr^\alpha$ for all $r \geq 0$ and some constant $C = C(V)$.

In other words for all $v \in \mathbb{R}^n$ with $\|v\| \leq 1$ and for all x, y in a relatively compact set V of U one has $||v|_x - |v|_y|| \leq C\|x - y\|^\alpha$ for some constant $C = C(V)$.

From the considerations above the following is clear:

Example 3.1. Let M be a $\mathcal{C}^{1,\alpha}$ manifold with a \mathcal{C}^α Finsler structure. If $\phi : N \rightarrow M$ is a $\mathcal{C}^{1,\alpha}$ immersion, then the induced Finsler structure on N is \mathcal{C}^α . In particular each $\mathcal{C}^{1,\alpha}$ submanifold of M inherits a \mathcal{C}^α Finsler structure.

The next example is immediately checked in a fixed chart:

Example 3.2. Let M be a $\mathcal{C}^{1,\alpha}$ manifold with a \mathcal{C}^α Finsler structure. Then for a locally α -Hölder function $f : M \rightarrow \mathbb{R}^+$ the Finsler structure $|\cdot|' := f|\cdot|$ on M given by $|v|'_x = f(x)|v|_x$ is again \mathcal{C}^α .

The last example is probably the most interesting one:

Example 3.3. Let M_1, M_2 be two manifolds with boundaries N_1, N_2 . Let g_1 resp. g_2 be Riemannian metrics on M_1 and on M_2 . Assume, that there is a $\mathcal{C}^{1,\alpha}$ diffeomorphism $\phi : N_1 \rightarrow N_2$ with $\phi^*(g_2) = g_1$. Consider the manifold M that arises from gluing M_1 and M_2 along the identifications of their boundaries given by the isometry ϕ . The manifold M is $\mathcal{C}^{1,\alpha}$ and the Riemannian metric g that coincides with g_1 on M_1 and with g_2 on M_2 is \mathcal{C}^α . It is interesting to note, that even if M, N_1, N_2 and ϕ are smooth the resulting metric g is not better than Lipschitz. It can be smooth only in very special cases. Usually it even does not have locally one-sided curvature bounds (compare [Kos02b] and [Kos02a]).

3.2 Comparison to the tangent spaces

Since all our results are of local nature we will restrict ourselves for the rest of the section to a chart $U \subset \mathbb{R}^n$, that is a convex subset of \mathbb{R}^n . So let $|\cdot|$ be a continuous Finsler structure on U . We can assume (considering a relatively compact subset of U instead of U) that the function $o_U(r) = o(r)$ as defined above is finite and bounded above by $C_1 > 2$ for all r . Moreover for all $x \in U$ we have $\frac{1}{C_2} \|\cdot\| \leq |\cdot|_x \leq C_2 \|\cdot\|$ for some constant $C_2 \geq 2$. Hence for the Finsler metric d and all $x, y \in U$ one has $\frac{1}{C_2} d(x, y) \leq \|x - y\| \leq C_2 d(x, y)$.

For a Lipschitz curve $\gamma : [a, b] \rightarrow U$ we will denote its Euclidean length by $L_\epsilon(\gamma)$. For each $x \in U$ we denote by $L^x(\gamma)$ the length of γ in the normed space $(\mathbb{R}^n, |\cdot|_x)$, i.e. $L^x(\gamma) = \int_a^b |\gamma'(t)|_x dt$. We will refer this length as the length relative to x . From our assumptions on U we see that $\frac{1}{C_2} L_\epsilon(\gamma) \leq L(\gamma) \leq C_2 L_\epsilon(\gamma)$.

The following lemma allows us to control the relative lengths of curves in terms of their actual lengths.

Lemma 3.1. *Let $\gamma : [0, s] \rightarrow U$ be a Lipschitz curve starting at $x = \gamma(0)$. Then we have $\|L(\gamma) - L^x(\gamma)\| \leq o(L_\epsilon(\gamma)) L_\epsilon(\gamma)$.*

Proof. We may assume that γ is parametrized by the Euclidean length. Then $L(\gamma) = \int_0^s |\gamma'(t)|_{\gamma(t)} dt = L^x(\gamma) + \int_0^s (|\gamma'(t)|_{\gamma(t)} - |\gamma'(t)|_x) dt$.

However by definition we get $\| |\gamma'(t)|_{\gamma(t)} - |\gamma'(t)|_x \| \leq o(\|x - \gamma(t)\|)$. Since $\|x - \gamma(t)\| \leq t$ and the function o is non-decreasing, we get $\|L(\gamma) - L^x(\gamma)\| \leq \int_0^s o(t) dt \leq o(s)s = o(L_\epsilon(\gamma)) L_\epsilon(\gamma)$. \square

Using the same argument of the proof of Lemma 3.1 one can also prove the following lemma, which allows to compare the relative lengths of a curve.

Lemma 3.2. *Let $\gamma : [0, s] \rightarrow U$ be a Lipschitz curve. $L^x(\gamma) - L^y(\gamma) \leq o(\|x - y\|) L_\epsilon(\gamma)$ for all $x, y \in U$.*

The following results are immediate applications of Lemma 3.1.

Corollary 3.3. *For all $x, y \in U$ one has $\|d(x, y) - |x - y|_x\| \leq o(\|x - y\|)\|x - y\|$, where d denotes the Finsler metric.*

In the case $o(r) \leq Cr^\alpha$ where $C = C(U)$ is a constant depending only U and $1 \geq \alpha > 0$ we obtain:

Corollary 3.4. *If the Finsler structure on U is C^α , then $\|L(\gamma) - L^x(\gamma)\| \leq C'L(\gamma)^{1+\alpha}$ and $\|d(x, z) - |x - z|_x\| \leq C'd(x, z)^{1+\alpha}$ for $C' = C_2^{\alpha+1}C$.*

4. THE SMOOTHNESS OF GEODESICS

Let M be a manifold with a C^α Finsler structure. We have already noticed that the statements are of the local nature. Therefore from now on we fix a chart U on M and assume that U be an open ball in \mathbb{R}^n with a C^α Finsler structure. We denote by d the Finsler metric and by $\|\cdot\|$ the Euclidean metric on U and $|\cdot|_x$ denotes the norm at point x induced from the Finsler structure. We assume that all the constants introduced in Section 3.2 are the same. Thus let $C > 2$ be a constant, such that for all $x \in U$ we have $\frac{1}{C}\|\cdot\| \leq |\cdot|_x \leq C\|\cdot\|$ and that the function o satisfies $o(r) = o_U(r) \leq Cr^\alpha$. We have $\frac{1}{C}d(x, y) \leq \|x - y\| \leq Cd(x, y)$ and $\|d(x, y) - |x - y|_x\| \leq \|x - y\|^{1+\alpha}$. Moreover we assume that for all x, y in U and for all Lipschitz curves γ in U starting at x one has $\|L^x(\gamma) - L(\gamma)\| \leq CL^x(\gamma)^{1+\alpha}$ and $\|L^x(\gamma) - L(\gamma)\| \leq CL(\gamma)^{1+\alpha}$. Also $\|d(x, y) - |x - y|_x\| \leq Cd(x, y)^{1+\alpha}$ and $\||x - y|_x - |x - y|_y\| \leq C\|x - y\|^{1+\alpha}$ for all $x, y \in U$.

4.1 The estimations of Hartmann and Calabi and proof of Theorem 1.3

Here we assume in addition to the above hypothesis that for each $x \in U$ the norm $|\cdot|_x$ is κ -convex of type p . Let $h < 1$ be a small positive number and let $\gamma : [-h, h] \rightarrow U$ be a geodesic. Set $x = \gamma(-h)$, $y = \gamma(0)$ and $z = \gamma(h)$. Let $\eta : [-h, h] \rightarrow U$ be the straight line between x and z parametrised proportional to the Euclidean arc length and put $m = \eta(0) = \frac{1}{2}(x + z)$. We will estimate the distance $|y - m|_x$.

Denote the geodesic sub-segment of γ connecting x and y by γ_1 . By Corollary 3.3 we have $|x - y|_x \leq L_x(\gamma_1) \leq L(\gamma_1) + CL(\gamma_1)^{1+\alpha} \leq h + Ch^{1+\alpha}$. Similarly denote $\gamma([0, h])$ by γ_2 . Thus $|y - z|_x \leq |y - z|_y + C\|y - z\| \|x - y\|^\alpha \leq L(\gamma_2) + CL(\gamma_2)^{1+\alpha} + C^2d(y, z) (Cd(x, y))^\alpha \leq h + (C + C^3)h^{1+\alpha}$. Moreover we have $|x - z|_x \geq d(x, z) - Cd(x, z)^{1+\alpha} \geq 2h - 4Ch^{1+\alpha}$.

Finally since $|\cdot|$ is κ -convex of type p applying Lemma 2.3 for $K = \max\{C + C^3, 4C\}$ we see that $|y - m|_x \leq 2\lambda h^{1+\frac{\alpha}{p}}$ for all $h \leq \epsilon$ where $\lambda = \lambda(K, \kappa)$ and $\epsilon = \epsilon(K, \kappa)$ are the constants given in Lemma 2.3. This means

exactly that $\|\gamma(-h) + \gamma(h) - 2\gamma(0)\| \leq 2\lambda h^{1+\frac{\alpha}{p}}$, and hence by Lemma 2.2 this implies Theorem 1.3.

4.2 Better estimations and proof of Theorem 1.4

Before to give the proof of Theorem 1.4 we make two trivial remarks that are used in the following arguments.

Remark 4.1. Let $f : [a, b] \rightarrow \mathbb{R}$ be a positive continuous function. Then $\|b - a\| \int_a^b f^2(t) dt \geq (\int_a^b f(t) dt)^2$.

Remark 4.2. There exists an ϵ (for example $\epsilon = \frac{1}{10}$) such that for all real numbers $x, a, b > 0$ with $\|1 - x\| + \|a\| + \|b\| < \epsilon$ one has $\sqrt{x + a - b} \geq \sqrt{x} + \frac{1}{3}a - b$. In fact this follows directly from Taylor expansion of the square root function.

Let U be a chart with the properties as defined in the beginning of the section and assume in addition, that all tangent spaces $(T_x U, |\cdot|_x)$ are Euclidean spaces, i.e. the norm $|\cdot|_x$ stems from the a scalar product $\langle \cdot, \cdot \rangle_x$. Therefore the norm is κ -convex of type 2 with $\kappa \geq \frac{1}{10}$ (see example 2.2).

Let again h be a small number, $\gamma : [-h, h] \rightarrow U$ a geodesic, $x = \gamma(-h)$, $z = \gamma(h)$, $y = \gamma(0)$ and $m = \frac{1}{2}(x + z)$. We may assume that m is the origin of \mathbb{R}^n and that the norm $|\cdot|_0$ is the Euclidean norm $\|\cdot\|$. Set $e = \frac{z}{\|z\|}$ and denote by P the line generated by e . Let H be the orthogonal hyperplane to P . Remark that for each unit vector $h \in H$ and all $a \in U$ we have $\langle e, h \rangle_a = \frac{1}{4}(|e + h|_a^2 - |e - h|_a^2) = (|e + h|_0^2 - |e + h|_a^2) + (|e - h|_0^2 - |e - h|_a^2)$. Since $\|v\|_x - \|v\|_y \leq C\|x - y\|^{\alpha+1}$ for all x, y and v with $\|v\| = 1$ we obtain $\langle e, h \rangle_a \leq 10C^2(\|e + h\| + \|e - h\|)\|a\|^\alpha \leq 10C^2\|a\|^\alpha$.

We decompose $\gamma(t)$ as $\gamma(t) = f(t)e + v(t)$, with $v(t) \in H$ and $f(t) \in \mathbb{R}$. By Theorem 1.3 the geodesics are uniformly locally $\mathcal{C}^{1, \frac{\alpha}{2}}$. Therefore f and v are of class \mathcal{C}^1 and by Lemma 2.1 choosing h small enough we may assume $\|v'(t)\| \leq \epsilon$ for arbitrary small $\epsilon > 0$. Since $|\gamma'(t)|_{\gamma(t)} = 1$ we see that $f'(t)|e|_{\gamma(t)} \geq 1 - \epsilon$.

Consider the straight line $\eta : [-h, h] \rightarrow U$ between x and z parametrised by $\eta(t) = f(t)e$. We see that $\gamma(t) = \eta(t) + v(t)$. Moreover $1 - f'(t)|e|_{\gamma(t)} \leq \epsilon$ and $\|v'(t)\| \leq \epsilon$ for arbitrary ϵ and we can use Remark 4.2, and obtain

$$\begin{aligned} 1 &= |\gamma'(t)|_{\gamma(t)} = (f'(t)^2 |e|_{\gamma(t)}^2 + |v'(t)|_{\gamma(t)}^2 + 2f'(t)\langle e, v'(t) \rangle_{\gamma(t)})^{\frac{1}{2}} \\ &\geq f'(t)|e|_{\gamma(t)} + \frac{1}{3}(|v'(t)|_{\gamma(t)}^2) - 2|f'(t)\langle e, v'(t) \rangle_{\gamma(t)}| \\ &\geq f'(t)|e|_{\eta(t)} - C\|v(t)\|^\alpha + \frac{1}{C'}\|v'(t)\|^2 - C'\|v'(t)\|(\|\gamma(t)\|)^\alpha \end{aligned}$$

where $C' = \max\{\frac{C^2}{3}, 40C\}$.

Moreover $L(\gamma) \leq L(\eta) = \int_{-h}^h f'(t)|e|_{\eta(t)} dt$, since γ is a geodesic. This together with the above inequality implies $\frac{1}{C'} \int_{-h}^h \|v'(t)\|^2 \leq C' \int_{-h}^h (\|v(t)\|^\alpha + \|v'(t)\| \|\gamma(t)\|^\alpha) dt$.

Using $\|\gamma(t)\| \leq \|v(t)\| + 2\|t\|$ and $\|v'(t)\| \leq 1$, we see that $\int_{-h}^h \|v'(t)\|^2 \leq \frac{1}{2}K(\int_{-h}^h (\|v(t)\|)^\alpha + \|v'(t)\| \|t\|^\alpha dt)$ where $K = 2C'^2$. Thus one has

$$\text{either } \int_{-h}^h \|v'(t)\|^2 dt \leq K \int_{-h}^h (\|v(t)\|)^\alpha dt \quad (1),$$

$$\text{or } \int_{-h}^h \|v'(t)\|^2 dt \leq K \int_{-h}^h \|v'(t)\| \|t\|^\alpha dt \quad (2).$$

Consider the maximum $v_0 = \max_{-h \leq t \leq h} \|v(t)\|$. Since $v(h) = v(-h) = 0$ we have $v_0 \leq \int_{-h}^h \|v'(t)\| dt$. In case (2) we get from the Cauchy Schwartz inequality $\int_{-h}^h \|v'(t)\|^2 dt \leq K(\int_{-h}^h \|v'(t)\|^2 dt)^{\frac{1}{2}}(\int_{-h}^h \|t\|^{2\alpha} dt)^{\frac{1}{2}}$. This implies $\int_{-h}^h \|v'(t)\|^2 dt \leq 2K^2 h^{2\alpha+1}$, and using Remark 4.1 we obtain $v_0 \leq 2K^2 h^{\alpha+1}$. In case (1), we obtain $(\int_{-h}^h \|v'(t)\| dt)^2 \leq 2Kh \int_{-h}^h (\|v(t)\|)^\alpha dt \leq 4Kh^2 v_0^\alpha$ using Remark 4.1. Therefore $v_0^2 \leq 4Kh^2 v_0^\alpha$. That implies $v_0 \leq K'h^{\frac{2}{2-\alpha}}$ where $K' = (4K)^{\frac{1}{2-\alpha}}$.

Since for $\beta = \frac{\alpha}{2-\alpha}$ we have $\beta < \alpha$ and $\frac{2}{2-\alpha} = 1 + \beta$, for each $t \in [-h, h]$ we obtain: $\|\gamma(t) - \eta(t)\| = C\|v(t)\| \leq Cv_0 \leq Ah^{1+\beta}$, for some A depending only on C .

In order to apply Lemma 2.2 we have to show that $\|\gamma(0) - m\| \leq Lh^{1+\beta}$ for some constant L depending only on C . Since m is the origin we have $\|\gamma(0) - m\| \leq \|\gamma(0) - \eta(0)\| + \|\eta(0)\| \leq Ah^{1+\beta} + \|f(0)e\|$. Thus we only have to show $\|f(0)e\| \leq Lh^{1+\beta}$ for some fixed L .

Assume $\|f(0)e\| \geq Lh^{1+\beta}$ with $L \geq 2C^2$. We may assume that $f(0) < 0$. We have $|x|_x = |x - m|_x \leq h + Ch^{1+\alpha}$. Thus $|x - \eta(0)|_x \leq h + Ch^{1+\alpha} - |f(0)e|_x \leq h - \frac{L}{2C}h^{1+\beta}$. Then $|x - \gamma(0)|_x \leq |x - \eta(0)|_x + |\eta(0) - \gamma(0)|_x \leq h - \frac{L}{2C}h^{1+\beta} + AC h^{1+\beta}$. If $L \geq 4AC^2$ we get $|x - \eta(0)|_x \leq h - \frac{L}{4C}h^{1+\beta}$. Using Corollary 3.4 we get a contradiction to $d(x, \gamma(0)) = h$. Thus we have prove $L \leq 4AC^2$ and the proof of Theorem 1.4 is finished.

5. THE COUNTEREXAMPLE

Let $0 < \alpha < 1$ be fixed and set $\beta = \frac{\alpha}{2-\alpha}$. Consider in \mathbb{R}^2 the graph Γ of the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(t) = |t|^{1+\beta}$. Set $\rho(x) = d(x, \Gamma) = \min_{y \in \Gamma} \{|x - y|\}$. Denote by \tilde{g} the Euclidean metric and consider on \mathbb{R}^2 the metric $g_x = (1 + A\rho(x)^\alpha)\tilde{g}$, where A is a constant ≥ 100 . This metric is α -Hölder and it is locally Lipschitz outside of Γ . At each point of Γ the new metric g coincides with the Euclidean metric \tilde{g} . For $t \in \mathbb{R}$ denote by x_t the point $(t, f(t))$ on Γ . We will prove that the geodesics between x_{-t} and x_t cannot be uniformly $\mathcal{C}^{1,l}$ for any $l > \beta$, i.e. we show that geodesic between x_{-t} and x_t must be very close to Γ .

The length of the curve $\gamma_t : [-t, t] \rightarrow \mathbb{R}^2$ defined by $\gamma_t(s) = (s, f(s)) = x_s$ is given by $L(\gamma_t) = L_e(\gamma_t) = \int_{-t}^t \sqrt{1 + (1 + \beta)^2 s^{2\beta}} ds$. Hence by Remark 4.2

and the fact that $\beta \leq 1$, we have for all sufficiently small t , $L(\gamma_t) \leq \int_{-t}^t (1 + 2s^{2\beta}) ds \leq 2t + 10t^{2\beta+1}$.

Now fix a small relatively compact neighbourhood U of 0 and assume that geodesics in U are uniformly $\mathcal{C}^{1,l}$ for some $l > \beta$. Let $\eta_t : [-h, h] \rightarrow \mathbb{R}^2$ be a geodesic between x_{-t} and x_t . Hence for all s, r in $[-t, t]$ we have $\|\eta'_t(s) - \eta'_t(r)\| \leq C\|s - r\|^l$ where C is a constant depending only on U . By Lemma 2.1 we see that for each $r \in [-h, h]$ the distance between $\eta_t(r)$ and the affine line between x_{-t} and x_t is at most Ct^{1+l} .

We reparametrise η_t by the first coordinate and denote the new curve by $\tilde{\eta}_t : [-t, t] \rightarrow \mathbb{R}^2$ where $\tilde{\eta}_t(s) = (s, y_t(s))$. From above we know $y_t(s) \geq f(t) - Ct^{l+1} = t^{\beta+1}(1 - Ct^{l-\beta})$. Since $l > \beta$ for arbitrary $\epsilon > 0$ and all t small enough we get $y_t(s) \geq f(t)(1 - \epsilon) \geq \frac{2}{3}f(t)$.

For each $x_r = (r, f(r)) \in \Gamma$ one has either $|r| \geq \frac{t}{2}$ or $f(r) \leq \frac{t^{1+\beta}}{2}$. Hence for each s with $|s| \leq \frac{t}{3}$ we get $d(\tilde{\eta}_t(s), \Gamma) \geq \frac{1}{6}t^{1+\beta}$.

Now we can estimate the length of η_t by $L(\eta_t) = L(\tilde{\eta}_t) \geq L_\epsilon(\tilde{\eta}_t) + \int_{-t}^t A(d(\tilde{\eta}_t(r), \Gamma))^\alpha dr \geq 2t + A \int_{-\frac{t}{3}}^{\frac{t}{3}} (\frac{1}{6}t^{1+\beta})^\alpha dr = 2t + \frac{100}{9}t^{1+\alpha+\beta}$.

But by our assumption η_t is a geodesic, hence $L(\eta_t) \leq L(\gamma_t)$. We get $10t^{2\beta+1} \geq \frac{100}{9}t^{1+\alpha+\beta}$. Since $1 + \alpha + \alpha\beta = 1 + 2\beta$ this is a contradiction.

REFERENCES

- [CH70] E. Calabi and Ph. Hartmann. On the smoothness of isometries. *Duke Math.J.*, 37:741–750, 1970.
- [Kos02a] Kosovskii. On gluing of Riemannian manifolds of curvature $\geq \kappa$. *Algebra i Analiz* 14, 14:140–157, 2002.
- [Kos02b] Kosovskii. On gluing of Riemannian manifolds of curvature $\leq \kappa$. *Algebra i Analiz*, 14:73–86, 2002.
- [LT79] J. Lindenstrauss and L. Tzafriri. *Classical Banach spaces. Function spaces*, volume 2. Springer-Verlag, 1979.
- [Lyt] A. Lytchak. Almost convex subsets. Preprint.
- [Tay] M. Taylor. Existence and regularity of Isometries. Preprint.